

Jgg93c_Hunstanton

Jones, G.G. 1993c The Animal Bones. In Bradley, R., Chowne, P., Cleal, R.M.J., Healy, F. and Kinnes, I., Excavations on Redgate Hill, Hunstanton, Norfolk, and at Tattershall Thorpe, Lincolnshire. *East Anglian Archaeology*, 57, 61-64, Microfiche Tables 36-7.

late neo EBA Hunstanton, Norfolk Jgg93c_Hunstanton

neo 305, EBA 35 ident. bones.

Late Neo 305 ident. C Sg P: 77, 16, 182, (30 other). Red 6 + 13a, roe 5, cat 2, dolphin 1, mouse (cf. *Apodemus*.) 3. Pig log ratio Fig.48.

EBA 35 ident. C Sg P 2, 25, 3+1sk, (4 other). Red 2 + 2a.

Below: Three extracts from Jgg Reports Summaries.accdb showing numbers;
an example original data page with method notes;
and the zones method description (from Jgg94a).

1List just first fields							
Id	pdf ref1	Site	Ph	Location	County	pdf ref2	Ph Ord
141	Jgg93c_Hunstanton	Hunstanton	LNeo	Norfolk	Norfolk	Jgg93c_Hunstanton	4.4
142	Jgg93c_Hunstanton	Hunstanton	EBA	Norfolk	Norfolk	Jgg93c_Hunstanton	5.2

2CSPH plus totals										
Id	pdf ref1	Ph	Cattle	Shgt	Pig	Horse	Other N	TotIdent	Total	Ph Ord
141	Jgg93c_Hunstanton	LNeo	77	16	182	0	30	305	1052	4.4
142	Jgg93c_Hunstanton	EBA	2	25	4	0	4	35		5.2

3Other species						
Id	pdf ref1	Ph	Other N	OtherM	Bird	Ph Ord
141	Jgg93c_Hunstanton	LNeo	30	red 6 + 13a, roe 5, cat 2, dolphin 1, mouse 3 (cf. <i>Apodemus</i>).		4.4
142	Jgg93c_Hunstanton	EBA	4	red 2+2a		5.2

See next pages.

An example original-record page from Aylesbury Prebendal.
Farley and Jones 2012.

ATP85 EIA

shy gt p.3

mb modern breakers.
ob old

hc R	1	-		
R				
skull	both	2	2	
L				
hc L	1	-		
mx R	1	-		
L	1	-		
md R	1	-		
L	2	-		
teeth				
HEAD 3				
atlas	1			
axis	3			
other cerv.	2			
thor	2	7		
lumb.	1	2		
sacr	1			
cdl	1			
st vt	3			
VT		5		
blade				
neck				
sc R	1	2		
L	1			
P	2			
Rs	2			
hu d	2	5		
P	2			
Ls	2			
d	2			
ra R	1			
d	1	3		
Ls	2			
d				
ul R	2	3		
Ls	2			
il	1			
R	1			
il	1			
epu	1			
pe	1	2		
is	1			
L	2			
is	1			
epu	1			
P				
Rs				
done				
straight				
onto				
sheet				
P				
s				
d				
fib/pat				
P				
Rs				
ti				
d				

ti	P		
Ls			
d			
ind. vs. Body		75	
calc R			
prox			
act L			
dist R			
L			
P			
Rs			
mc d			
P			
Ls			
d			
P			
Rs			
mt d			
P			
Ls			
d			
c/s dmp			
py-ab, mp			
ph lat		4	
2nd		3	
3rd		3	
ab. ph.			
FOOT		14	
ZONE			
TOTAL		92	
TOTAL		97	
CONTEXT			
MN			

3047
3048
3050
3053
see also 3059 Report
notes many - 2 pairs photos
A 3028 - all one, except col

3054
*
not all def. related

3055
3057
3058

Complete bone with all 6 zones:

- 1: prox epiphysis fused,
- 2: prox metaphysis fused,
- 3,4: upper shaft, lower shaft,
- 5: distal metaphysis fused,
- 6: distal epiphysis fused.

goat (gt) radius.
μ - Measured.
Radius not fused to ulna.

See ZonesJgg

A nearly complete femur, prox unfused metaphysis, upper and lower shaft, dist unfused metaphysis and distal loose epiphysis. No prox epiphysis.

The Zones method is described here, in:
Jones, G.G. 1994a Animal Bones. In Ayers, B.,
Excavations at Fishergate, Norwich, 1985.
East Anglian Archaeology, 68, 37.

II. Mammal and Bird Bone

by Gillian Jones

The mammal and bird bone from Fishergate, of late ninth century to late medieval date, is summarised in Table 7.

Method (Fig. 22)


The main bone assemblage was hand collected. A small quantity of bone was recovered from the sieved samples. Bone was recorded on two lists, with the more complete bones on a zone list and the other bones on a fragments list. On the zone list were recorded complete bones or bone pieces as follows:

Skull:

substantial pieces of horncore, frontal, lacrimal, malar, parietal, squamous temporal, occipital; upper jaw and mandible with at least one tooth present; loose teeth.

Long-bones:

where more than half of any of the six areas shown on Figure 22 was present and where the following small areas of bone were present: humerus, the distal posterior part of the shaft; radius, the proximal part of the ulnar groove; femur, the supracondylar fossa; tibia, the anterior, distal part of zone 4.



Division of Long Bones into six zones (figure after Baker and Brothwell, 1984 p.44).

Figure 22 Division of long-bones into six zones.

Other bones:

more than half the following bone or bone elements: vertebra, the body and central arch; scapula, the neck and glenoid cavity; ulna, the olecranon and proximal articulation; pelvis, the iliac shaft and the iliac, ischial and pubic parts of the acetabulum; calcaneum, the proximal part and articulation; the patella, astragalus and phalanx.

With cattle, substantial pieces of the ends of long-bones, even when less-than-half complete, were included on the zone list. This was done in order to avoid loss of important epiphyseal fusion data. However, few bones fell into this category, due to the well-preserved and relatively unfragmented nature of the bone assemblage.

The separation of the fragments in bone recording may be useful, in that it is likely to be less repeatable than that of the more-complete segment. Accurate identification of fragments probably varies somewhat between bone analysts, and for a single analyst depending on the time available for study. It will also tend to vary according to the number of similar-sized species present. Some fragments may be assigned to cattle which, if red deer and horse were as common as cattle, would have remained as 'large unidentified'. However, a fragment was not identified unless it bore clear features typical of the particular species.

Table 9 (microfiche), the Anatomical Analysis, shows the total number of bones (BN) and a reduced number of zones. For long-bones, these are zones 2 and 5, labelled p (proximal) and d (distal), and zone 4 for the humerus, femur and tibia, and zone 3 for the radius and metapodials, labelled s (shaft).

Dating

The dating of the bone is based on the identified site periods (see Chronological summary, p.ix) which were themselves dated by artefacts. There was, however, residual earlier pottery in later phases and some of the bone may therefore also be residual earlier material.

General description of the bone

The bone from the Period I marsh deposits was well-preserved and dominated by cattle. Many of the bones were fairly complete and had surfaces which were dark in colour and hard with little abrasion. The good state of preservation of the bone suggests that the marsh was used as a primary dump. In general few bones appeared to relate to each other. Upper and lower jaws of cattle from context 129 probably belong to each other, but, for example, no distal tibiae with matching astragalus were found and only two immature cattle bones were recovered as both metaphysis and epiphysis (against fourteen unfused metaphyses without epiphyses and eight epiphyses without metaphyses). Of thirty immature vertebral centra, in only one case was a matching epiphysis preserved.

Bone from Periods III1 and III2 was also well-preserved. The bone was less dark in colour than the Period I bone and some of it bore a sandy accretion. Again, few bones related to each other (upper and lower jaws, hock joint bones, or metaphyses and epiphyses). One might suggest that casual dumping of bone took place over time and that there may have been some post-depositional movement of bone in the deposits.

It is expected that access to the marsh to dump bone would favour the large bones of cattle and that the high percentage may be more informative about the particular area of the town than the general supply of meat in Nor-